

Are Damped Ly α Systems Rotating Disks ?

Cédric Ledoux¹, Patrick Petitjean², Jacqueline Bergeron³

¹*Observatoire Astronomique de Strasbourg, 11 Rue de l'Université, F-67000 Strasbourg, France*

²*Institut d'Astrophysique de Paris, CNRS, 98bis Boulevard Arago, F-75014 Paris, France*

³*European Southern Observatory, Karl-Schwarzschild-Straße 2, D-85748 Garching bei München, Germany*

Abstract. We report on high spectral resolution observations of five damped Ly α systems whose line velocity profiles and abundances are analyzed. By combining these data with information from the literature, we study the kinematics of the low and high ionization phases of damped systems and discuss the possibility that part of the motions is due to rotation.

1 The Si II λ 1808 and Fe II profiles

We study the general properties of the kinematics of the low and high ionization phases in a large sample of 26 damped Ly α systems (hereafter DLAS) covering the redshift range 1.17–4.38. It has been argued that high-redshift DLAS could represent a population of large, fast rotating disks mainly on the basis of the asymmetric shape of their low ionization species' absorption lines [3][2]. We note a strong correlation between the velocity broadenings of the Si II λ 1808 and Fe II λ 1608 transition lines for apparent optical depths of 0.1 and 0.7 respectively (see Fig. 1, left panel). This shows that the physical conditions are quite homogeneous in the neutral gas and that large variations of abundance ratios and thus large depletions onto dust grains in DLAS are unlikely. Since the Fe II transitions are easier to detect, we are able to compare velocities measured on *the same ion*.

2 Velocity broadening and asymmetry

We define an asymmetry parameter $\eta = |V_1 + V_2| / (V_1 - V_2)$ that is close to zero for a symmetric profile and close to one for a one-sided profile. It can be seen on Fig. 1 (right panel) that the Fe II line asymmetry is correlated with the velocity broadening up to $\Delta V \sim 120 \text{ km s}^{-1}$. For larger velocity broadenings however this correlation disappears. The systems with large Δv , such as the system at $z_{abs} = 3.151$ toward Q 2233+131, are indeed very peculiar, with kinematics much more consistent with random motions than ordered disks. They show sub-systems as those expected if the objects are in the process of




Figure 1: Velocity broadening of the Fe II $\lambda 1608$ lines at $\tau_\nu \sim 0.7$ versus that of Si II $\lambda 1808$ at $\tau_\nu \sim 0.1$ (left panel) and line asymmetry versus the velocity broadening for Fe II $\lambda 1608$ measured at $\tau_\nu \sim 0.7$ (right panel).

merging. Therefore the claim that CDM models should be rejected has to be considered with caution. Indeed using hydro-simulations, Haehnelt et al. (1997) [1] have reached the same conclusion. Although more work has to be done especially investigating comparisons with simple kinematical models, the overall picture seems to be more in favor of the DLAS to be aggregates of dense knots with complex kinematics rather than ordered disks. This does not rule out however the possibility that part of the kinematics could be due to rotation. The correlation shown in Fig. 1 (right panel) is suggestive of rotational motions in sub-systems on scales smaller than $\Delta V < 120 \text{ km s}^{-1}$. The kinematics of the low and high ionization species themselves is found to be statistically correlated though the high ionization phase has a much more disturbed kinematical field than low ions. This should be taken into account in any model of high-redshift damped Ly α systems.

Acknowledgements. The participation of C. Ledoux to the conference has been funded by the E.C.

References

- [1] Haehnelt M.G., Steinmetz M., Rauch M., 1997, astro-ph/9706201
- [2] Prochaska J.X., Wolfe A.M., 1997, astro-ph/9704169
- [3] Wolfe A.M., 1995, eds Meylan G., in *QSO Absorption Lines*. Springer Editions, Berlin, p. 13